ELSEVIER

Toxicology 127 (1998) 243–244

TOXICOLOGY

Volume contents

Volume 127 (1998)

Repeated dosing with the peroxisome proliferator clofibrate decreases the toxicity of model hepatotoxic agents in male mice
Protection was afforded against 24 hour toxicity of bromobenzene, carbon tetrachloride or chlorofrom
J.E. Manautou, V.M. Silva, G.E. Hennig, H.E. Whiteley (USA) 1–10

Immunoenzymatic determination of atrazine in rat tissue samples
Quantitation demonstrates comparatively low tissue levels of atrazine
B. Scutaru, T. Giersch, C. Cozmei, B. Hock (Romania, Germany) 11–16

The influence of polymorphonuclear leukocytes on altered pulmonary epithelial permeability during ozone exposure
Excessive accumulation of leukocytes alone does not increase permeability
P.G. Reinhart, D.J. Bassett, D.K. Bhalla (USA) 17–28

Subchronic effects of smokeless tobacco extract (STE) on hepatic lipid peroxidation, DNA damage and excretion of urinary metabolites in rats
Administration of STE every other day for 105 days caused multiple signs of oxidative stress
M. Bagchi, D. Bagchi, E.A. Hassoun, S.J. Stohs (USA) 29–38

Formic acid excretion in rats exposed to trichloroethylene: a possible explanation for renal toxicity in long-term studies
Formic acid may be the toxic intermediate
T. Green, J. Dow, J.R. Foster, P.M. Hext (UK) 39–47

Evaluation of P19 cells for studying mechanisms of developmental toxicity: application to four direct-acting alkylating agents
The established cell line is suitable for mechanistic studies in developmental toxicology
M.R. Seeley, E.M. Faustman (USA) 49–58

Aluminum transport out of brain extracellular fluid is proton dependent and inhibited by mersalyl acid, suggesting mediation by the monocarboxylate transporter (MCT1)
A proton-dependent process appears to be involved in the removal of aluminum from brain extracellular fluid
D.C. Ackley, R.A. Yokel (USA) 59–67

Urinary and biliary excretion kinetics of 1-hydroxypyrene following intravenous and oral administration of pyrene in rats
Urinary and biliary kinetics of 1-hydroxypyrene were similar regardless of dose and exposure route
M. Bouchard, C. Viau (Canada) 69–84

US EPA's IRIS pilot program: establishing IRIS as a centralized, peer-reviewed data base with agency consensus
Review of EPS's IRIS (Integrated Risk Information System) and its Pilot Program
A. Mills, G.L. Foureman (USA) 85–95

Structural specificities and significance for coeliac disease of wheat gliadin peptides able to agglutinate or to prevent agglutination of K562(S) cells
About a dozen gliadin peptides were synthesized and tested for the aminoacid sequences producing agglutination
M. De Vincenzi, A. Stammati, R. Luchetti, M. Silano, G. Gasbarrini, V. Silano (Italy) 97–106

Stressor-specific enhancement of hsp induction by low doses of stressors in conditions of self- and cross-sensitization
The mutual influences of low dose arsenite and cadmium as well as heat are described
F.A. Wiegant, N. Spieker, R. van Wijk (The Netherlands) 107–119

Effects of molybdate on the sulfation of harmol and α-naphthol
Molybdate may inhibit the sulfation of highly sulfated compounds
J.W. Boles, C.D. Klaassen (USA) 121–127
Ability of 16 priority PAHs to be directly cytotoxic to a cell line from the rainbow trout gill
Naphthalene was the only compound cytotoxic below its solubility limit
K. Schirmer, D.G. Dixon, B.M. Greenberg, N.C. Bols (Canada) 129–141
Ability of 16 priority PAHs to be photocytotoxic to a cell line from the rainbow trout gill
Fluoranthene and pyrene appear to have the most phototoxic potential
K. Schirmer, A.G. Chan, B.M. Greenberg, D.G. Dixon, N.C. Bols (Canada) 143–155
Induction of pulmonary fibrosis in organ-cultured rat lung by cadmium chloride and transforming growth factor-β1
There is an additive or synergistic action of both factors
C.-J. Lin, P.-C. Yang, M.-T. Hsu, F.-H. Yew, T.-Y. Liu, C.-T. Shun, S.-W. Tyan, T.-C. Lee (Taiwan, ROC) 157–166
Role of placental metallothionein in maternal to fetal transfer of cadmium in genetically altered mice
Placental metallothionein is mostly located in the cytoplasm of spongiotrophoblasts, glycogen cells and yolk sac
J.C. Lau, M.G. Joseph, M.G. Cherian (Canada) 167–178
PCB congener 77-induced ultrastructural alterations in the rat liver: a quantitative study
Increased volume of smooth endoplasmic reticulum and of mitochondria was observed
C. Gilroy, B.J. Connell, A. Singh, P. Suidgeest, I. Chu (Canada) 179–185
Urinary excretion levels of sodium and potassium in environmental cadmium-exposed subjects
Increased K excretion is a more sensitive indicator of Cd exposure than is Na excretion
K. Monzawa, T. Kido, H. Yamaya, E. Kobayashi, K. Nogawa (Japan) 187–193
Induction of apoptosis with fusarenon-X in mouse thymocytes
The trichotecene mycotoxin FX induces apoptosis in thymocytes
K. Miura, Y. Nakajima, N. Yamanaka, K. Terao, T. Shibato, S. Ishino (Japan) 195–206
Side effects of brequinar and brequinar analogues, in combination with cyclosporine, in the rat
Brequinar and analogues potentiate the toxicity of cyclosporin
C. Pally, D. Smith, B. Jaffee, R. Magolda, H. Zehender, B. Dorobek, P. Donatsch, C. Papageorgiou, H.-J. Schuurman (Switzerland, USA) 207–222
Acute and subchronic immunotoxicity of *p*-chloronitrobenzene in mice. I. Effect on natural killer, cytotoxic T-lymphocyte activities and mitogen-stimulated lymphocyte proliferation
Parachloronitrobenzene decreased NK cell and cytotoxic T lymphocyte activity and LPS-induced lymphocyte proliferation in mice
Q. Li, M. Minami, H. Inagaki (Japan) 223–232

Announcements 233–235
Author index 237
Subject index 239–241
Volume contents 243–244